arXiv:1405.4458v1 [math.GT] 18 May 2014

SINGULAR MEASURES ON THE LIMIT SET OF A KLEINIAN GROUP

WOOJIN JEON

Abstract. We consider a finitely generated torsion free Kleinian group H and a random walk on H with respect to a symmetric nondegenerate probability measure μ with finite support. When H is geometrically infinite without parabolics or when H is Gromov hyperbolic with parabolics, we prove that the Patterson-Sullivan measure is singular with respect to the harmonic measure coming from μ.

1. Introduction

Let H be a finitely generated *Kleinian group*, i.e., a discrete subgroup of $\mathrm{PSL}(2,\mathbb{C})$. We assume that H has no elliptic elements. Since $\mathrm{PSL}(2,\mathbb{C})$ is the group of orientation preserving isometries of the hyperbolic 3-space $\mathbb{H}^3$, we can consider the action of H on $\mathbb{H}^3$ which can be continuously extended to the ideal boundary S^2_∞ of $\mathbb{H}^3$. The *limit set* Λ_H of H is the set of limit points of an orbit $H\cdot o$ where $o\in\mathbb{H}^3$. This definition does not depend on the choice of o and Λ_H is contained in S^2_∞.

We consider two measures on Λ_H. One is the Patterson-Sullivan measure ρ_x based at $x\in\mathbb{H}^3$ and the other is the harmonic measure ν_H coming from a random walk on H. Given a probability measure μ on H and the corresponding random walk $\{Y_n\}$ on H, we can describe ν_H as follows. For $A\subset S^2_\infty$, $\nu_H(A)$ is the probability for $Y_n\cdot o$ to converge to a point in $A\subset S^2_\infty$. When H is a *Fuchsian group* i.e., a discrete subgroup of $\mathrm{PSL}(2,\mathbb{R})$, H acts isometrically on $\mathbb{H}^2$ and Y. Guivarc'h and Y. Le Jan proved the following.

Theorem 1.1. *(Y. Guivarc'h and Y. Le Jan* [16]*) Let H is a discrete subgroup of* $\mathrm{PSL}(2,\mathbb{R})$ *such that $\mathbb{H}^2/H$ is a noncompact surface with finite area. Let ν_H is the harmonic measure on S^1_∞ coming from a symmetric nondegenerate probability measure μ with finite support on H. Then ν_H is singular with respect to the Lebesgue measure on S^1_∞.*

In this paper, we generalize Theorem 1.1 to Kleinian groups as follows.

[1]School of Mathematics KIAS, Hoegiro 87, Dongdaemun-gu, Seoul, 130-722, Korea
email : jwoojin@kias.re.kr

[2]2010 *Mathematics Subject Classification.* 51M10, 57S25, 60J50

[3]*Key words and phrases.* Kleinian group, Cannon-Thurston map, Poisson boundary, harmonic measure, Patterson-Sullivan measure

Theorem 1.2. *Suppose that H is a finitely generated Kleinian group such that*

(1) H is Gromov hyperbolic.
(2) H is not convex-cocompact.
(3) The orbit map $\tau_o : H \to \mathbb{H}^3$ sending h to $h \cdot o$ extends continuously to $\widehat{\tau}_o : \partial H \to S^2_\infty$.

Here ∂H is the Gromov boundary of H. Let ν_H be the harmonic measure on Λ_H coming from a symmetric nondegenerate probability measure μ with finite support on H. Then ν_H is singular with respect to the Patterson-Sullivan measure on Λ_H.

We remark that when $\Lambda_H = S^2_\infty$, the Patterson-Sullivan measure ρ_o where o is the origin of the Poincaré ball model is equal to the Lebesgue measure $Leb_{S^2_\infty}$ on S^2_∞ up to constant multiple. Thus every Patterson-Sullivan measure ρ_x is equal to $Leb_{S^2_\infty}$ up to homothety. The continuous extension $\widehat{\tau}_o$ of the orbit map τ_o for H is called the *Cannon-Thurston map* of H and its existence has been verified recently for surface Kleinian groups by [24]. The paper [25] dealt with the general case. For the proof of Theorem 1.2, we modify and extend the argument used in the proof of [3, Prop 5.4, Prop 5.5]. The assumption of the existence of the Cannon-Thurston map is crucial.

2. Preliminaries

2.1. **Hyperbolic spaces.** For $\delta \geq 0$, a geodesic metric space (X, d) is called *δ-hyperbolic* if for any geodesic triangle in X, each side of the triangle is contained in the δ-neighborhood of the union of the other two sides. We call a geodesic metric space a *hyperbolic space* (in the sense of Gromov) if it is δ-hyperbolic for some $\delta \geq 0$. A hyperbolic metric space (X, d) has a boundary at infinity ∂X called the *Gromov boundary* which can be defined as follows. We say a sequence $\{x_n\}$ in (X, d) *converges to infinity* if

$$\liminf_{i,j \to \infty} (x_i | x_j)_x = \infty$$

for some (hence every) basepoint x, where

$$(y|z)_x := \frac{1}{2}(d(x, y) + d(x, z) - d(y, z))$$

for $x, y, z \in X$. This product $(y|z)_x$ is called the *Gromov product* of y and z with respect to x and approximates within 2δ the distance from x to any geodesic $[y, z]$ joining y, z. Two sequences $\{x_n\}$, $\{y_n\} \subset X$ converging to infinity are said to be *equivalent* if

$$\liminf_{i,j \to \infty} (x_i | y_j)_x = \infty$$

for some $x \in X$. The Gromov boundary ∂X is defined as the set of the equivalence classes of sequences converging to infinity in X. This definition is independent of the choice of the base point $x \in X$. The Gromov product

can be extended to ∂X. We can define a natural metric $d_\epsilon(\cdot,\cdot)$ on ∂X such that $C^{-1}e^{-\epsilon(p|q)_x} \leq d_\epsilon(p,q) \leq Ce^{-\epsilon(p|q)_x}$ where $(p,q) \in \partial X \times \partial X$.

Fixing a finite generating set S of H, the word metric $d_w(\cdot,\cdot)$ on H is defined by setting $d_w(g,h) = |g^{-1}h|_S$ where $|g|_S$ is the minimum of the number of elements of S whose product is g. When we consider the right Cayley graph Γ_H of H with the length of every edge being 1, $d_w(g,h)$ is the minimum of the lengths of paths joining g and h in Γ_H. A finitely generated group H is called as a *hyperbolic group* if its Cayley graph Γ_H is hyperbolic with respect to the word metric for a finite generating set S of H. We simply denote the word length of g as $|g|$ by fixing a finite generating set. We denote the Gromov boundary of Γ_H by ∂H. For basic properties of Gromov hyperbolic spaces, hyperbolic groups and their Gromov boundaries, we refer the reader to [15, 13] and [20].

2.2. **Kleinian groups.** Let H be a finitely generated *Kleinian group*. We assume that H has no elliptic elements. The *limit set* Λ_H of H is the set of limit points of an orbit $H \cdot o$ where $o \in \mathbb{H}^3$. The complementary open set $\Omega_H = S^2_\infty \backslash \Lambda_H$ becomes the domain of discontinuity of the action of H and by Ahlfors' finiteness theorem, Ω_H/H consists of finitely many Riemann surfaces which are called as the *conformal boundaries at infinity* of H.

The *convex hull* of H is defined to be the smallest convex set in $\mathbb{H}^3$ whose closure in $\mathbb{H}^3 \cup S^2_\infty$ containes Λ_H. The *convex core* of H is the quotient of its convex hull by H itself. The Kleinian group H is called *geometrically finite(resp. convex cocompact)* if its convex core has finite volume(resp. if its convex core is compact). Every convex cocompact Kleinian group is a hyperbolic group by the Schwarz-Milnor lemma(See [5] for example). By Thurston's uniformization theorem in [28], every geometrically infinite Kleinian group without parabolics has a convex cocompact representation which is faithful. Thus if H is geometrically infinite without parabolics, then it is hyperbolic.

If the orbit map $\tau_o : H \to \mathbb{H}^3$ sending $h \to h \cdot o$ can be extended continuously to $\widehat{\tau}_o : \partial H \to S^2_\infty$, we call $\widehat{\tau}_o$ as the *Cannon-Thurston map* of H following [9, 4, 22, 23, 24, 25, 27]. If H is convex cocompact, then τ_o is a quasi-isometric embedding and we have a natural homeomorphism $\widehat{\tau}_o$ from the Gromov boundary ∂H to the limit set Λ_H. When H is a geometrically finite Kleinian group with parabolics, we still have a continuous extension of τ_o from the Floyd boundary (See [12, 14]) of H to Λ_H. If we further assume H is hyperbolic, then the Floyd boundary of H is equal to the Gromov boundary ∂H and a parabolic element h gives two points $\{h_\infty, h_{-\infty}\}$ in ∂H such that $\widehat{\tau}_o(h_\infty) = \widehat{\tau}_o(h_{-\infty})$ where $\{h_\infty, h_{-\infty}\}$ are the accumulation points of $\{h^i | i \in \mathbb{Z}\}$ in Γ_H.

Now we consider the case when H is a geometrically infinite hyperbolic Kleinian group and we assume the Cannon-Thurston map $\widehat{\tau}_o$ exists. In this case, we can find an exiting sequence of closed geodesics $\{c_n\}$ in $\mathbb{H}^3/H$. The exiting property of $\{c_n\}$ gives a bi-infinite quasigeodesic l in Γ_H such that

the two endpoints of l are identified by $\widehat{\tau}_o$. See [26] for details. We need the following lemma for the proof of Theorem 1.2.

Lemma 2.1. *Let H be a finitely generated Kleinian group with $\mathbb{H}^3/H$ not being convex cocompact and such that H is Gromov hyperbolic. We assume that the Cannon-Thurston map $\widehat{\tau}_o : \partial H \to S^2_\infty$ exists. Then there exists $\{h_n\} \subset H$ such that for any constant $D > 0$, $|h_n| - Dd_{\mathbb{H}^3}(o, h_n \cdot o) \to \infty$.*

Proof. We fix a finite generating set S of H. If H has a parabolic element g, then we can just take h_n as g^n. If not, then H is geometrically infinite and there exists a bi-infinite quasigeodesic l in Γ_H joining two points p, q in ∂H such that $\widehat{\tau}_o(p) = \widehat{\tau}_o(q)$. We represent l as a bi-infinite sequence of vertices in Γ_H

$$\{\cdots g_{-n}, g_{-(n-1)}, \cdots, id, \cdots, g_n, g_{n+1}, \cdots\}$$

such that $|g_i^{-1} g_{i+1}| = 1$. Then since $\widehat{\tau}_o(p) = \widehat{\tau}_o(q)$ and $\widehat{\tau}_o$ is a continuous extension of τ_o, we have $(g_{-n} \cdot o | g_n \cdot o)_o \to \infty$. If $\tau_o|_l$ is a quasi-isometric embedding then by the Morse-Mostow lemma(See [5] for example), the image of $\tau_o|_l$ is contained in a uniform neighborhood of a bi-infinite geodesic in Γ_H. Since $(g_{-n} \cdot o | g_n \cdot o)_o \to \infty$, this is not possible and thus $\tau_o|_l$ cannot be a quasi-isometric embedding. We note that $d_{\mathbb{H}^3}(g, h) \leq C|g^{-1}h|$ by the triangle inequality where $C := \max_{s \in S} d_{\mathbb{H}^3}(o, s \cdot o)$. We can also see the following. There exists a sequence of integers $i(n)$ and $j(i) > i$ such that for any $D > 0$, $|g_i^{-1} g_j| - Dd_{\mathbb{H}^3}(g_i \cdot o, g_j \cdot o) \to \infty$ as $n \to \infty$. We take h_n as $g_{i(n)}^{-1} g_{j(i)}$. □

2.3. **Poisson boundary.** Let μ be a probability measure on a group H.

- μ is called *nondegenerate* if the support of μ generates H as a semigroup.
- μ is called *symmetric* if $\mu(A) = \mu(A^{-1})$ where $A \subset H$ and A^{-1} is the set of the inverses of elements of A.

The word length $|h|$ of an element h in H is with respect to a fixed finite generating set of H. The *first moment* of (H, μ) is defined to be $\sum_{h \in H} |h| \mu(h)$. The *entropy* of (H, μ) is $\lim_{n \to \infty} - \sum_{h \in H} \mu(h) \log \mu(h)$. When μ has a finite support, both of its entropy and its first moment are finite.

The *random walk* on H with respect to μ is a Markov chain $\{Y_n\}$ with the transition probabilities given by $p_{h_1, h_2} = \mu(h_1^{-1} h_2)$. We denote $\{Y_n\}$ by Y. A random walk Y can also be described as a sequence of independent random variables $\{Z_n\}$ with values in the probability space (H, μ) such that $Y_n = Z_0 Z_1 \cdots Z_n$. When $Y_0 = id$, the random walk starts from the identity element id of H. We regard $Z = \{Z_n\}$ as an element of the product probability space $(H^{\mathbb{N}}, \mu^{\mathbb{N}})$ and Y as an element of the Kolmogrov representation space $(H^{\mathbb{N}}, \mathbb{P}_{id})$ where

$$\mathbb{P}_{id}(\{Y_n\} | Y_0 = id, Y_1 = h_1, Y_2 = h_2, \cdots, Y_n = h_n\}) = p_{id, h_1} p_{h_1, h_2} \cdots p_{h_{n-1}, h_n}$$

and $\mathbb{P}_{id}(\{Y_n\}|Y_0 \neq id\})$ is defined to be zero. We denote $\mathbb{P}_{id}$ by $\mathbb{P}$.

The time shift operator T acts on $(H^{\mathbb{N}}, \mathbb{P})$ by $(TY)_n = Y_{n+1}$ and defines an equivalence relation $\sim$ by saying $Y \sim Y'$ if and only if there exist positive integers k, k' such that $T^k Y = T^{k'} Y'$.

Definition 2.2. *The Poisson boundary of H with respect to μ is the quotient space of $(H^{\mathbb{N}}, \mathbb{P})$ by the smallest measurable equivalence relation generated by the equivalence relation $\sim$.*

The smallest measurable equivalence relation generated by $\sim$ is called as the *measurable envelope* [31]. When H is a hyperbolic group, we can consider the hitting measure ν of a random walk Y on ∂H and the Poisson boundary of H can be identified with $(\partial H, \nu)$ by [17, 18]. More precisely, it can be shown that $\{Y_n\}$ converges to a point in ∂H for almost every sample path $\{Y_n\}$ and $\nu(A)$ can be defined as the probability for a sample path $\{Y_n\}$ converges to a point in $A \subset \partial H$. Thus ν can also be called as *the law of* Y_∞. When H is a hyperbolic Kleinian group, H acts on $\mathbb{H}^3$ and we can consider the law of $Y_\infty \cdot o$ if $\{Y_n \cdot o\}$ converges to a point in S^2_∞ for almost every sample path $\{Y_n\}$. In fact, the law of $Y_\infty \cdot o$ is the push forward of ν by $\widehat{\tau}_o$ by the following Theorem 2.3 which summarizes [21, Corollary 6.2].

Theorem 2.3. *(Karlsson and Margulis* [21]*) Let H be a countable group isometrically acting on a uniformly convex, complete metric space (X, d) which is nonpositively curved in the sense of Busemann. Let μ be a probability measure on H with $\sum_{h \in H} d(o, h \cdot o)\mu(h) < \infty$. Let ∂X be the ideal boundary of X consisting of asymptotic classes of geodesic rays. Then*

- *Almost every sample path $\{Z_n\}$ in $H^{\mathbb{N}}$ with respect to $\mu^{\mathbb{N}}$ converges to a geodesic ray in X.*

Thus we have a map $\xi : (H^{\mathbb{N}}, \mu^{\mathbb{N}}) \to \partial X$ and we give ∂X the push-forward measure $\xi_*(\mu^{\mathbb{N}})$ so that ξ is measurable.

- *$(\partial X, \xi_*(\mu^{\mathbb{N}}))$ becomes the Poisson boundary of (H, μ) if the lattice counting function is subexponential, i.e., if*

$$\#\{h \in H | d(o, h \cdot o) < r\} \leq e^{Cr}$$

for some constant $C > 0$.

A similar form of Theorem 2.3 was also mentioned in Remark 3 following [18, Theorem 7.7]. It is known that the lattice counting function is subexponential for a large class of discrete subgroups of the isometry groups of Cartan-Hadamard manifolds. In particular, for pinched negatively curved case, it is proven in [34, Theorem 3.6.1]. Thus every Kleinian group has a subexponential lattice counting function. Moreover, since we are assuming H is finitely generated, we have $d_{\mathbb{H}^3}(o, h \cdot o) \leq D \cdot |h|$ by the triangle inequality.

Therefore we can apply Theorem 2.3 to a Kleinian group H acting on $\mathbb{H}^3$. We assume μ is a symmetric, nondegenerate probability measure on H with

finite support. Let ν_H be the law of $Y_\infty \cdot o$. Then (S^2_∞, ν_H) becomes the Poisson boundary of (H, μ) and thus $\widehat{\tau}_o : (\partial H, \nu) \to (S^2_\infty, \nu_H)$ is a measurable isomorphism.

2.4. **Conformal density.** For x in $\mathbb{H}^3$, the *Busemann function* $b_{x,\eta}(\cdot)$ at η with $b_{x,\eta}(x) = 0$ can be defined by choosing a geodesic ray $\alpha(t)$ from $\alpha(0) = x \in \mathbb{H}^3$ toward $\eta \in S^2_\infty$ as follows.

$$b_{x,\eta}(y) = \lim_{t\to\infty} (d_{\mathbb{H}^3}(y, \alpha(t)) - d_{\mathbb{H}^3}(x, \alpha(t)))$$

Lemma 2.4. *(*[29, Lemma 3.2.1]*) Consider the Poincaré ball model* $\{x \in \mathbb{R}^3 : |x| < 1\}$ *of* $\mathbb{H}^3$ *where* $|x|$ *is the usual Euclidean norm of* x*. Then*

$$e^{b_{x,\eta}(y)} = \frac{P(x,\eta)}{P(y,\eta)}$$

where $P(x, \eta)$ *is the Poisson kernel* $(1 - |x|^2)/|x - \eta|^2$*. Thus* $b_{x,\eta}(y)$ *is a continuous function of* $x, y \in \mathbb{H}^3$ *and* $\eta \in S^2_\infty$*.*

Fixing x, y in $\mathbb{H}^3$, the Poincaré series for H is

$$g_s(x, y) = \sum_{h\in H} e^{-s d_{\mathbb{H}^3}(x, h\cdot y)}$$

The critical exponent δ_H of H is defined as

$$\delta_H = \limsup_{r\to\infty} \frac{1}{r} \log(\#\{h \in H | d^3_{\mathbb{H}}(o, h \cdot o) \le r\})$$

Equivalently, δ_H is the infimum of the set of s such that $g_s(x, y)$ is finite. We call H is *divergent* if $g_s(x, y)$ diverges at $s = \delta_H$. Otherwise H is called *convergent*. These definitions are independent of the choices of $x, y \in \mathbb{H}^3$. Consider a family of measures $\{\rho^s_x\}$ defined by

$$\rho^s_x = \frac{1}{g_s(y, y)} \sum_{h\in H} e^{-s d_{\mathbb{H}^3}(x, h\cdot y)} \delta_{h\cdot y}$$

where $\delta_{h\cdot y}$ is the Dirac measure at $h \cdot y$. Then we can find a sequence $\{s_i\}$ with $s_i \to \delta^-_H$ such that $\rho^{s_i}_x$ weakly converges to a finite measure ρ_x on the compact space $\mathbb{H}^3 \cup S^2_\infty$. When H is divergent, ρ_x has its support on the limit set Λ_H. Since $\{\rho_x\}$ satisfies

$$\frac{d\rho_x}{d\rho_y}(\eta) = e^{-\delta_H b_{x,\eta}(y)}, h^*\rho_x = \rho_{h^{-1}\cdot x}$$

it becomes an H-invariant *conformal density* of dimension δ_H. Here the pull-back measure $h^*\rho_x$ is defined by setting $(h^*\rho_x)(A) := \rho_x(h \cdot A)$ for $A \subset \Lambda_H$. Even when H is convergent, we still can construct a conformal density by increasing the Dirac mass on each orbit point suitably [30, 32] and we denote a resulting conformal density also by ρ_x. We call ρ_x as the *Patterson-Sullivan measure* with base point $x \in \mathbb{H}^3$ in either case of H being convergent or divergent.

As an application of the Tameness theorem(see section 9 in [6] and [11, Prop. 3.9]), a Kleinian group H is divergent if and only if either H is geometrically finite or $\Lambda_H = S^2_\infty$. When $\Lambda_H = S^2_\infty$, ρ_x is equal to $Leb_{S^2_\infty}$ up to homothety and the diagonal action of H on $S^2_\infty \times S^2_\infty \backslash \Delta$ is ergodic with respect to $Leb_{S^2_\infty} \otimes Leb_{S^2_\infty}$. Here Δ means the diagonal set.

2.5. **Harmonic Density.** The *Green function* on a group H with a probability measure μ is defined by

$$G(g,h) = \sum_{n=0}^{\infty} \mu^n(g^{-1}h)$$

where μ^n is the n-th convolution power of μ. Then $F(g,h) = G(g,h)/G(h,h)$ is the probability that there exists $n \in \mathbb{N}$ such that $gY_n = h$. The Green metric d_G is defined by

$$d_G(g,h) = -\log F(g,h)$$

If H is a hyperbolic group and if μ is finitely supported nondegenerate symmetric probability measure on H, then d_G is a left invariant hyperbolic metric quasi-isometric to the word metric on H by [3, Corollary 1.2].

The *Martin kernel* $K : H \times H \to \mathbb{R}$ is defined by

$$K(g,h) = \frac{F(g,h)}{F(id,h)}$$

There exist constants $\{C_g\}$ such that for all $h \in H$, $K(g,h) \leq C_g$ for each $g \in H$. Let $H \cup \partial_M H$ be the metric completion of H with respect to the metric d_M defined by

$$d_M(h_1,h_2) = \sum_{h \in H} D_h \frac{|K(h,h_1) - K(h,h_2)| + |\delta_{h,h_1} - \delta_{h,h_2}|}{C_h + 1}$$

where $\{D_h\}$ is chosen so that $\sum_{h \in H} D_h < \infty$. $\delta_{h,g}$ is the Kronecker delta. Then it can be shown that $H \cup \partial_M H$ is in fact a compactification of H and the Martin kernel can be continuously extended to $H \times (H \cup \partial_M H)$. We call $\partial_M H$ as the *Martin boundary* of H. When H is a hyperbolic group, $\partial_M H$ is homeomorphic to the Gromov boundary of H. For this, see [2, 19] or [3, Corollary 1.8].

A positive function u on H is called *harmonic* if $u(h) = \sum_{g \in H} \mu(h^{-1}g)u(g)$ for all $h \in H$ and the Martin representation theorem says for any harmonic function u on H, there exists a measure ν_u on $\partial_M H$ such that

$$u(h) = \int_{\partial_M H} K(h,\xi) d\nu_u(\xi)$$

When we take u as a constant function $u \equiv 1$, the support of ν_u with the measure $\nu_u = \nu_1$ becomes the Poisson boundary of H. It is known that almost every sample path $\{Y_n\}$ converges to a point on $\partial_M H$. Thus for

a hyperbolic group H, we can identify ν with ν_1. We have the following change of variable formula.

$$\frac{d(h^*\nu)}{d\nu}(\xi) = K(h^{-1}, \xi)$$

For a hyperbolic group H, we can define the Busemann function on H with respect to the Green metric d_G by

$$b^G_{id,\xi}(h) = \sup_{\{h_n\}} \limsup_{n\to\infty} (d_G(h_n, id) - d_G(h_n, h))$$

where sup is taken over all sequences $\{h_n\}$ converging to ξ. If we choose $\{h_n\}$ along a quasigeodesic ray toward ξ in Γ_H, then we can just take the usual limit as $h_n \to \xi$ to define $b^G_{id,\xi}(h)$. By the definition of the Martin kernel, we get

$$\frac{d(h^*\nu)}{d\nu}(\xi) = e^{b^G_{id,\xi}(h^{-1})}$$

We have the same formula for the push-forward measure ν_H on S^2_∞ which was used in the proof of [3, Prop. 5.5]

Lemma 2.5. *Let H be a Kleinian group which is hyperbolic and let ν_H be the push-forward measure $\widehat{\tau}_o(\nu)$ on S^2_∞ as before. Then*

$$\frac{d(h^*\nu_H)}{d\nu_H}(\widehat{\tau}_o(\xi)) = e^{b^G_{id,\xi}(h^{-1})}$$

where $\xi \in \partial H$ and $(h^*\nu_H)(A) = \nu_H(h\cdot A)$ for $A \subset S^2_\infty$ for any ν_H-measurable set $A \subset S^2_\infty$.

Now we apply [19, Theorem 3.3] to the Cayley graph of the hyperbolic group H. Since the Gromov boundary ∂H itself is the conical limit set of H, we can see that the diagonal action of H on $\partial H \times \partial H \backslash \Delta$ with respect to the measure class of $\nu \otimes \nu$ is ergodic.

Lemma 2.6. *Let H be a Kleinian group which is hyperbolic. Let μ be a symmetric nondegenerate probability measure on H with finite support. Then the diagonal action of H on $S^2_\infty \times S^2_\infty \backslash \Delta$ is ergodic with respect to $\nu_H \otimes \nu_H$.*

Proof. We know $(\partial H, \nu)$ and (S^2_∞, ν_H) are measurably isomorphic by the equivariant map $\widehat{\tau}_o$ by Theorem 2.3. □

3. Singularity of measures on Λ_H

In this section, we prove Theorem 1.2. Recall that for $\eta_1, \eta_2 \in S^2_\infty$, the *Busemann cocycle* $B_o(\eta_1, \eta_2)$ is defined as $b_{o,\eta_1}(y) + b_{o,\eta_2}(y)$ where y is any point in the bi-infinite geodesic l joining η_1 and η_2. Geometrically $B_o(\eta_1, \eta_2)$ is the length of the geodesic subsegment of l contained in the intersection of the horoballs passing through o and centered at η_1, η_2. Thus it is nonnegative for any $(\eta_1, \eta_2) \in S^2_\infty \times S^2_\infty \backslash \Delta$ and it is zero if and only if o is contained in l.

Proof of Theorem 1.2. We follow and extend the proof of [3, Prop. 5.4, Prop. 5.5]. Define the measure $\widetilde{\rho}_o$ on $S^2_\infty \times S^2_\infty \backslash \Delta$ by

$$d\widetilde{\rho}_o(\eta_1, \eta_2) = e^{2\delta_H B_o(\eta_1, \eta_2)} d\rho_o(\eta_1) d\rho_o(\eta_2)$$

Then $\widetilde{\rho}_o$ is a H-invariant measure although it may not be an ergodic measure with respect to the H-action. We define the measure $\widetilde{\nu}_H$ on $S^2_\infty \times S^2_\infty \backslash \Delta$ as

$$d\widetilde{\nu}_H(\widehat{\tau}_o\xi_1, \widehat{\tau}_o\xi_2) = e^{2(\xi_1|\xi_2)^G_{id}} d\nu_H(\widehat{\tau}_o\xi_1) d\nu_H(\widehat{\tau}_o\xi_2)$$

Then $\widetilde{\nu}_H$ is invariant under the action of H by [19, Prop. 2.2]. Furthermore it is an ergodic measure by Corollary 2.6. Now we suppose ν_H is equivalent to ρ_o and we claim that the Radon-Nikodym derivative $d\rho_o/d\nu_H$ is ν_H-essentially upper and lower bounded by positive constants.
Proof of the claim: We have a ν_H-integrable function J and a $\widetilde{\nu}_H$-measurable function $\widetilde{J}$ defined by $d\nu_H = J d\rho_o$ and $d\widetilde{\nu}_H = \widetilde{J} d\widetilde{\rho}_o$. Since $\widetilde{J}$ is positive almost everywhere, there exists a constant $C > 0$ such that the set $A := \{(\eta_1, \eta_2) \in S^2_\infty \times S^2_\infty \backslash \Delta | C^{-1} \leq \widetilde{J}(\eta_1, \eta_2) \leq C\}$ has positive $\widetilde{\nu}_H$-measure. Since $\widetilde{\nu}_H$ is ergodic, there exists $h \in H$ such that $(h\eta_1, h\eta_2) \in A$ for $\widetilde{\nu}_H$-almost every (η_1, η_2). Since $\widetilde{\rho}_o$ is also H-invariant, we have $\widetilde{J}(\eta_1, \eta_2) = \widetilde{J}(h\eta_1, h\eta_2)$ and thus A has the full $\widetilde{\nu}_H$-measure. When $\eta_i = \widehat{\tau}_o\xi_i$ for $i = 1, 2$, we have

$$\widetilde{J}(\eta_1, \eta_2) = J(\eta_1) J(\eta_2) \frac{e^{2(\xi_1|\xi_2)^G_{id}}}{e^{2\delta_H B_o(\eta_1, \eta_2)}}$$

Thus we have for $\widetilde{\nu}_H$-almost every (η_1, η_2),

$$C^{-1} \frac{e^{2\delta_H B_o(\eta_1, \eta_2)}}{e^{2(\xi_1|\xi_2)^G_{id}}} \leq J(\eta_1) J(\eta_2) \leq C \frac{e^{2\delta_H B_o(\eta_1, \eta_2)}}{e^{2(\xi_1|\xi_2)^G_{id}}}$$

Assume, seeking a contradiction, J is unbounded in $B_s \cap \Lambda_H \subset S^2_\infty$ where B_s is a small spherical open ball in S^2_∞. We choose another small spherical open ball B'_s far from B_s so that for all $(\eta, \eta') \in B_s \times B'_s$, $D^{-1} < e^{B_o(\eta, \eta')} < D$ for some constant $D > 0$. Since $\widehat{\tau}_o$ is a continuous map, the distance in ∂H from the open set $\widehat{\tau}_o^{-1}(B'_s)$ to the open set $\widehat{\tau}_o^{-1}(B_s)$ has a positive lower bound. If we let $\widehat{\tau}_o\xi = \eta$ and $\widehat{\tau}_o\xi' = \eta'$, this means $(\xi'|\xi)_{id}$ has an upper bound and the same is true for $(\xi'|\xi)^G_{id}$. Thus $e^{(\xi'|\xi)^G_{id}}$ is positively upper and lower bounded. From this, we get that $J(\eta)J(\eta')$ is positively upper and lower bounded for $\widetilde{\nu}_H$-almost all $(\eta, \eta') \in B_s \times B'_s$. But there is a constant $D_1 > 0$ such that $B := \{\eta' \in B'_s | D_1^{-1} \leq J(\eta') \leq D_1\}$ has a positive measure so that $B_s \times B$ has a positive $\widetilde{\nu}_H$-measure. Since J is unbounded on B_s, $J(\eta)J(\eta')$ cannot be essentially bounded in $B_s \times B$. This is a contradiction and we have proved our original claim.

Now we recall the change of variable formulas for conformal and harmonic measures.

$$\frac{d(h^*\rho_o)}{d\rho_o}(\eta) = e^{\delta_H b_{o,\eta}(h^{-1}\cdot o)}$$

$$\frac{d(h^*\nu_H)}{d\nu_H}(\widehat{\tau}_o(\xi)) = e^{b^G_{id,\xi}(h^{-1})}$$

Since the density of ρ_o with respect to ν_H is uniformly bounded away from zero, there exists $C_1 > 0$

$$|\operatorname*{ess\,sup}_{\eta\in\Lambda_H} \delta_H b_{o,\eta}(h^{-1}\cdot o) - \operatorname*{ess\,sup}_{\xi\in\partial H} b^G_{id,\xi}(h^{-1})| < C_1$$

By [3, Lemma 2.5], there exists a constant $C_2 > 0$ such that

$$|d_G(id, h^{-1}) - \operatorname*{ess\,sup}_{\xi\in\partial H} b^G_{id,\xi}(h^{-1})| < C_2$$

Here we can replace 'esssup' by 'sup' because every open set in ∂H has a positive ν-measure and $b^G_{id,\xi}(h^{-1})$ is a continuous function with respect to ξ by Lemma 2.4. We also have

$$\sup_{\eta\in\Lambda_H} b_{o,\eta}(h^{-1}\cdot o) \le d_{\mathbb{H}^3}(o, h^{-1}\cdot o)$$

by the triangle inequality. Therefore there exist C_3(which may not be a positive number) such that for all $h \in H$,

$$\delta_H d_{\mathbb{H}^3}(o, h^{-1}\cdot o) - d_G(id, h^{-1}) > C_3$$

But by Lemma 2.1, there exists a sequence $\{h_n\} \subset H$ such that $|h_n^{-1}| - \delta_H d_{\mathbb{H}^3}(o, h_n^{-1}\cdot o)$ goes to infinity. Since the word metric is quasi-isometric to the Green metric, we have a contradiction. □

Note that even in the case of $\Lambda_H = S^2_\infty$, Theorem 1.2 is not a direct consequence of [3, Prop. 4.5]. In fact, [3, Prop. 4.5] is using [3, Prop. 4.4] which is valid for the case that τ_o is a quasi-isometry and $\widehat{\tau}_o$ is the natural homeomorphism between ∂H and S^2_∞. For the case that τ_o is not a quasi-isometry, we need to assume the existence of the continuous boundary extension $\widehat{\tau}_o$ of τ_o.

References

[1] I. Agol, *Tameness of hyperbolic 3-manifolds*, preprint, arXiv:math.GT/0405568 (2004)

[2] A. Ancona, *Théorie du potentiel sur les graphes et les variétés*, École d'été de Probabilités de Saint-Flour XVIII-1988, Lecture Notes in Math. 1427, Springer (1990), 1-112.

[3] S. Blachère, P. Haïssinsky and P. Mathieu, *Harmonic measures versus quasiconformal measures for hyperbolic groups*, Ann. Sci. Éc. Norm. Supér. (4) 44 (2011), 683-721.

[4] B. H. Bowditch, *The Cannon-Thurston map for punctured surface groups*, Math. Z. 255 (2007), 35-76.

[5] M. R. Bridson, A. Haefliger, *Metric spaces of non-positive curvature*, Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 319 (1999) Springer-Verlag, Berlin.

[6] D. Canary, *Ends of hyperbolic 3-manifolds*, Journal of Amer. Math. Soc., Vol 6. no. 1 (1993), 1-35.

[7] A. J. Casson, S. A. Bleiler, Automorphisms of surfaces after Nielsen and Thurston, London Math. Soc. Student Texts. 9. (1988).

[8] D. Calegari and D. Gabai, *Shrinkwrapping and the taming of hyperbolic 3-maifolds*, J. Amer. Math. Soc. 19 (2006), no.2, 385-446.

[9] J. Cannon and W. P. Thurston, *Group invariant Peano Curves*, Geometry and Topology 11 (2007), 1315-1356.
[10] M. Coornaert, *Mesures de Patterson-Sullivan sur le bord d'un espace hyperbolique au sens de Gromov*, Pac. J. Math. 159, no. 2 (1993), 241-270.
[11] M. Culller, P. Shalen, *Paradoxical decompositions, 2-generators Kleinian groups, and volumes of hyperbolic 3-manifolds*, J. Amer. Math. Soc 5 (1992), no. 2, 231-288.
[12] W. Floyd, *Group completions and limit sets of Kleinian groups.* Invent. Math. 57 (1980), 205 -218.
[13] E. Ghys and P. de la Harpe, editors. *Sur les Groups Hyperbolic d'après Mikhael Gromov*, vol. 83 of Progress in Mathematics. Birkhäuser, (1990).
[14] V. Gerasimov, *Floyd maps for relatively hyperbolic groups*, GAFA 22 (2012), 1-39.
[15] M. Gromov, *Hyperbolic groups*, In S. M. Gersten, editor, Essays in group theory, volume 8 of MSRI publications (1987), 75-263. Springer-Verlag.
[16] Y. Guivarch and Y. Le Jan, *Sur l'enroulement du flot géodésique*, C. R. Acad. Sci. Paris Sér. I Math. 311 (1990), 645-648.
[17] V. A. Kaimanovich, *Poisson boundaries of hyperbolic groups*, C. R. Ac. Sci. Paris, Ser. I 318 (1994), 59-64.
[18] V. A. Kaimanovich, *The Poisson formula for groups with hyperbolic properties*, Ann. of Math. 152 (2000), 659-692.
[19] V. A. Kaimanovich, *Ergodicity of harmonic invariant measures for the geodesic flow on hyperbolic spaces*, Journal für die reine und angewandte Mathematik 455 (1994), 57-104.
[20] I. Kapovich and N. Benakli, *Boundaries of hyperbolic groups*, In Combinatorial and geometric group theory (New York, 2000/Hoboken, NJ, 2001), volume 296 of Contemp. Math., 39-93. Amer. Math. Soc., (2002) Providence, RI.
[21] A. Karlsson and G. A. Margulis, *A multiplicative ergodic theorem and nonpositively curved spaces*, Commun. Math. Phy. 208 (1999), 107-123.
[22] C. McMullen, *Local connectivity, Kleinian groups and geodesics on the blow-up of the torus*, Invent. math. 97 (2000), 95-127.
[23] Y. N. Minsky, *On rigidity, limit sets and end invariants of hyperbolic 3-manifolds*, J. Amer. Math. Soc. 7 (1994), 539-588.
[24] M. Mj, *Cannon-Thurston map for Surface groups*, preprint, arXiv:math.GT/0511041 (2006), To appear in Ann. of Math.
[25] M. Mj, *Cannon-Thurston maps for Kleinian groups*, preprint, arXiv:1002.0996.
[26] M. Mj, *Ending laminations and Cannon-Thurston maps*, arXiv:math 0701725 (2007), To appear in Geometric and Functional analysis.
[27] M. Mj, *Cannon-Thurston maps for pared manifolds of bounded geometry*, Geometry and Topology 13 (2009), 189-245.
[28] J. Morgan, *On Thurston's uniformization theorem for three dimensional manifolds, The Smith Conjecture*, Academic Press (1984), 37-125.
[29] P. J. Nicholls, *The ergodic theory of discrete groups*, London Math. Soc. Lecture Note Series 143 (1989), Cambridge Univ. Press.
[30] S. J. Patterson, *The limit set of a fuchsian group*, Acta math. 136 (1976), 241-273.
[31] V. A. Rokhlin, *On the fundamental ideas of measure theory*, Amer. Math. Soc. Trans. 71 (1962).
[32] D. Sullivan, *The density at infinity of a discrete group of hyperbolic motions*, Publ. math. IHES 50 (1979), 171-202.
[33] W. P. Thurston, *The geometry and topology of 3-manifolds*, Lecture note (1980), available on http://library.msri.org/nonmsri/gt3m/.
[34] C. Yue, *The ergodic theory of discrete isometry groups on manifolds of variable negative curvature*, Trans. Amer. Math. Soc. Volume 348, no.12 (1996), 4965-5005.

Woojin Jeon
School of Mathematics
KIAS, Hoegiro 87, Dongdaemun-gu
Seoul, 130-722, Korea
jwoojin@kias.re.kr